08963656 Results

Antibody SEQ ID NO: 2

Result No.	Score	Query Match	Length	DB	ID	Description
1	355	100.0	355	19	AAW51744	Human C-C chemokin
2	308	86.8	355	17	AAW03376	CC-chemokine recep
3	308	86.8	355	18	AAW10100	Human C-C chemokin
4	228	64.2	355	17	AAW03377	CC-chemokine recep
5	228	64.2	355	18	AAW31850	Human eosinophil e
6	228	64.2	355	18	AAW27124	Human chemokine re
7	228	64.2	355	19	AAW51745	Human C-C chemokin
8	228	64.2	355	22	AAG80109	Human CCR3 protein
9	228	64.2	356	18	AAW25943	Human CCKR3 chemok
10	190	53.5	355	22	ABB56341	Non-endogenous hum
11	134	37.7	355	17	AAW03378	CC-chemokine recep
12	134	37.7	355	19	AAW51746	Human C-C chemokin
13	41	11.5	295	22	AAG80106	Human CCR1 protein
14	41	11.5	355	15	AAR52749	C-C chemokine rece
15	41	11.5	355	18	AAW26588	Human MIP-1 alpha/
16	41	11.5	355	18	AAW25751	Human MIP-lalpha/R
17	41	11.5	355	21	AAB20571	Human CC-chemokine

```
RESULT 15
AAW26588
ID
    AAW26588 standard; Protein; 355 AA.
XX
AC
     AAW26588;
XX
\mathtt{DT}
     21-JAN-1998 (first entry)
ХX
DE
     Human MIP-1 alpha/RANTES receptor.
XX
KW
     Macrophage inflammatory protein-1 alpha; MIP-1 alpha;
KW
     reduced upon activation normal T expressed and secreted; RANTES;
KW
     receptor; cytokine; antiinflammatory; inflammation; human.
XX
os
     Homo sapiens.
XX
PN
     US5652133-A.
\mathbf{X}\mathbf{X}
PD
     29-JUL-1997.
xx
PF
     28-JAN-1993;
                     93US-0012988.
\mathbf{x}\mathbf{x}
PR
     28-JAN-1993;
                    93US-0012988.
XX
PA
     (USSH ) US DEPT HEALTH & HUMAN SERVICES.
XX
ΡI
     Murphy PM;
ХX
DR
     WPI; 1997-392945/36.
     N-PSDB; AAT90384.
DR
xx
PT
     MIP-1-alpha and RANTES receptor nucleic acid - used to develop
PT
     products for the detection of these cytokine(s) and their receptors,
PT
     particularly in inflammatory processes
XX
PS
     Claim 2; Column 15-18; 12pp; English.
XX
CC
     This polypeptide comprises a claimed receptor for human macrophage
CC
     inflammatory protein-1 alpha (MIP-1 alpha) and regulated upon
CC
     activation normal T expressed and secreted (RANTES) protein. Also
CC
     claimed are: a nucleic acid (see AAT90384) that encodes the receptor;
     a subsequence of the nucleic acid, having at least 12 contiguous
```

```
CC
    nucleotides; a cell transformed or transfected with the nucleic
    acid; and purified MIP-1 alpha/RANTES receptor polypeptide. The
    products can be used for detecting the MIP-1 alpha/RANTES receptor
CC
CC
    and polymorphisms in physiological samples. In addition, the
    receptor can be expressed and used to assay for MIP-la/RANTES in
CC
CC
    biological samples. The quantitation of MIP-1 alpha/RANTES is
CC
    useful for monitoring the levels of these cytokines in a patient.
CC
    Such measurements are useful in following the antiinflammatory
CC
    effects of drugs and prospective usefulness of new antiinflammatory
CC
XX
so
    Sequence 355 AA;
                        11.5%; Score 41; DB 18; Length 355;
 Query Match
 Best Local Similarity 100.0%; Pred. No. 1.4e-31;
                                               0; Indels
          41; Conservative
                            0; Mismatches
                                                           0: Gaps
                                                                        0:
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Oν
         Db
     115 tglyseiffiilltidrylaivhavfalrartvtfgvitsi 155
```

```
8
Result
               Ouerv
  No.
        Score Match Length DB ID
                                                         Description
       . - - - - - - - - - - - -
          228 64.2
                        355 4 US-08-575-967A-4
                                                          Sequence 4, Appli
    2
          228 64.2
                        355 4 US-08-847-296B-1
                                                          Sequence 1, Appli
                       355 4 US-09-045-583-54
355 1 US-08-012-988A-2
    3
          228
               64.2
                                                          Sequence 54, Appl
               11.5
    4
           41
                                                          Sequence 2, Appli
               11.5
                        355 1 US-08-450-393A-5
                                                          Sequence 5, Appli
                       355 4 US-08-446-669-5
    6
           41
               11.5
                                                          Sequence 5, Appli
    7
           41
                11.5
                        355 4 US-09-045-583-53
                                                          Sequence 53, Appl
                       355 4 US-09-239-938-1
    8
           41
               11.5
                                                          Sequence 1, Appli
    9
           41
                11.5
                        355 5 PCT-US95-00476-5
                                                          Sequence 5, Appli
                        31 1 US-08-450-393A-14
31 4 US-08-446-669-14
   10
           31
                8.7
                                                          Sequence 14, Appl
   11
           31
                8.7
                                                          Sequence 14, Appl
                        31 5 PCT-US95-00476-14
                                                          Sequence 14, Appl
   12
           31
                8.7
   13
           22
                        31 1 US-08-450-393A-13
                6.2
                                                          Sequence 13, Appl
```

```
RESULT 15
PCT-US95-00476-13
; Sequence 13, Application PC/TUS9500476
  GENERAL INFORMATION:
    APPLICANT: The Regents of the University of California
    TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT
    TITLE OF INVENTION: PROTEIN RECEPTORS
    NUMBER OF SEQUENCES: 14
    CORRESPONDENCE ADDRESS:
      ADDRESSEE: Robbins, Berliner & Carson
      STREET: 201 N. Figueroa Street, 5th Floor
      CITY: Los Angeles
      STATE: California
      COUNTRY: USA
      ZIP: 90012-2628
    COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
      OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
      APPLICATION NUMBER: PCT/US95/00476
      FILING DATE:
      CLASSIFICATION:
    ATTORNEY/AGENT INFORMATION:
      NAME: Berliner, Robert
```

```
REGISTRATION NUMBER: 20,121
;
      REFERENCE/DOCKET NUMBER: 5555-291
    TELECOMMUNICATION INFORMATION:
      TELEPHONE: 310-977-1001
      TELEFAX: 310-977-1003
      TELEX:
  INFORMATION FOR SEQ ID NO: 13:
    SEQUENCE CHARACTERISTICS:
     LENGTH: 31 amino acids
      TYPE: amino acid
      TOPOLOGY: linear
    MOLECULE TYPE: peptide
PCT-US95-00476-13
                         6.2%; Score 22; DB 5; Length 31;
 Query Match
 Best Local Similarity 100.0%; Pred. No. 1e-13;
 Matches 22; Conservative 0; Mismatches
                                               0; Indels
                                                              0; Gaps
     121 IFFIILLTIDRYLAIVHAVFAL 142
         1 IFFIILLTIDRYLAIVHAVFAL 22
RESULT 13
US-08-450-393A-13
; Sequence 13, Application US/08450393A
: Patent No. 5707815
; GENERAL INFORMATION:
    APPLICANT: Charo, Israel
    APPLICANT: Coughlin, Shaun
    TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT TITLE OF INVENTION: PROTEIN RECEPTORS
    NUMBER OF SEQUENCES: 14
    CORRESPONDENCE ADDRESS:
      ADDRESSEE: Cooley Godward Castro Huddleson & Tatum
      STREET: 5 Palo Alto Square
      CITY: Palo Alto
STATE: California
      COUNTRY: USA
      ZIP: 94306-2155
    COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
      OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
     APPLICATION NUMBER: US/08/450,393A
      FILING DATE: May 25, 1995
      CLASSIFICATION: 424
    ATTORNEY/AGENT INFORMATION:
      NAME: Cserr, Luann
      REGISTRATION NUMBER: 31,822
      REFERENCE/DOCKET NUMBER: UCAL-237/02US
    TELECOMMUNICATION INFORMATION:
      TELEPHONE: 415-843-5165
      TELEFAX: 415-8857-0663
      TELEX: 380816CooleyPA
  INFORMATION FOR SEQ ID NO: 13:
    SEQUENCE CHARACTERISTICS:
      LENGTH: 31 amino acids
      TYPE: amino acid
     TOPOLOGY: linear
    MOLECULE TYPE: peptide
US-08-450-393A-13
                         6.2%; Score 22; DB 1; Length 31;
 Best Local Similarity 100.0%; Pred. No. 1e-13;
 Matches 22; Conservative 0; Mismatches 0; Indels
                                                             0; Gaps
```

```
RESULT 8
US-09-239-938-1
; Sequence 1, Application US/09239938
; Patent No. 6329510
; GENERAL INFORMATION:
; APPLICANT: Qin, Shixin
; APPLICANT: Newman, Walter
; APPLICANT: Kassam, Nasim
; APPLICANT: LeukoSite, Inc.
; TITLE OF INVENTION: ANTI-CCR1 ANTIBODIES AND METHODS OF USE
; TITLE OF INVENTION: THEREFOR
; FILE REFERENCE: LKS97-13
; CURRENT APPLICATION NUMBER: US/09/239,938
; CURRENT FILING DATE: 1999-01-29
; NUMBER OF SEQ ID NOS: 1
; SOFTWARE: FastSEQ for Windows Version 3.0
; SEQ ID NO 1
   LENGTH: 355
   TYPE: PRT
   ORGANISM: Homo sapien
US-09-239-938-1
 Query Match 11.5%; Score 41; DB 4; Length 355; Best Local Similarity 100.0%; Pred. No. 1.4e-30;
                                                 0; Indels
                                                               0; Gaps
 Matches 41; Conservative
                              0; Mismatches
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qу
          115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Db
```

121 IFFIILLTIDRYLAIVHAVFAL 142

Qу

DЪ

		₹				
Result		Query				
No.	Score	Match	Length	DB	ID	Description
1	308	86.8	355	2	G02436	chemokine (C-C) re
2	41	11.5	355	2	A45177	chemokine (C-C) re
3	33	9.3	355	2	149339	macrophage inflamm
4	33	9.3	359	2	I49341	MIP-1 alpha recept
5	22	6.2	360	2	JC2443	chemokine (C-C) re
6	22	6.2	374	2	I38450	chemokine (C-C) re
7	16	4.5	383	2	S55594	G protein-coupled
8	14	3.9	352	2	A43113	chemokine (C-C) re
9	12	3.4	356	2	I49340	MIP-1 alpha recept
10	12	3.4	360	2	A57160	chemokine (C-C) re
11	12	3.4	360	2	JC4587	chemokine (C-C) re
12	10	2.8	308	2	I50241	G protein-coupled
						<u>-</u>

```
RESULT 9
149340
MIP-1 alpha receptor like-1 - mouse
C;Species: Mus musculus (house mouse)
C;Date: 02-Jul-1996 #sequence_revision 02-Jul-1996 #text_change 13-Aug-1999
C;Accession: 149340
R;Gao, J.L.; Murphy, P.M.
J. Biol. Chem. 270, 17494-17501, 1995
A;Title: Cloning and differential tissue-specific expression of three mouse beta chemokine receptor-like genes, including the gene for a functional macrophage inflammatory protein-1 alpha receptor.
```

```
A; Reference number: 149339; MUID: 95340546
A; Accession: I49340
A; Status: preliminary; translated from GB/EMBL/DDBJ
A; Molecule type: DNA
A; Residues: 1-356 < RES>
A;Cross-references: EMBL:U28405; NID:g881549; PIDN:AAA89154.1; PID:g881550
C; Superfamily: vertebrate rhodopsin
  Query Match 3.4%; Score 12; DB 2; Length 356; Best Local Similarity 100.0%; Pred. No. 0.00064;
                                 0; Mismatches
                                                 0; Indels
                                                                  0: Gaps
                                                                              0:
  Matches 12; Conservative
      139 VFALRARTVTFG 150
Οv
          Db
      140 VFALRARTVTFG 151
RESULT 10
A57160
chemokine (C-C) receptor 4 - human
N; Alternate names: C-C CKR-4
C; Species: Homo sapiens (man)
C;Date: 10-Nov-1995 #sequence revision 10-Nov-1995 #text change 21-Jul-2000
C:Accession: A57160
R; Power, C.A.; Meyer, A.; Nemeth, K.; Bacon, K.B.; Hoogewerf, A.J.; Proudfoot, A.E.I.;
Wells, T.N.C.
J. Biol. Chem. 270, 19495-19500, 1995
A; Title: Molecular cloning and functional expression of a novel CC chemokine receptor
cDNA from a human basophilic cell line.
A; Reference number: A57160; MUID: 95370289
A; Accession: A57160
A; Status: preliminary; not compared with conceptual translation
A; Molecule type: mRNA
A; Residues: 1-360 < POW>
A;Cross-references: GB:X85740; NID:g1370103; PIDN:CAA59743.1; PID:g971452
A; Note: source clone K5-5
C; Genetics:
A; Gene: GDB: CMKBR4
A; Cross-references: GDB:677463
A; Map position: 3p21-3p21
C; Superfamily: vertebrate rhodopsin
C; Keywords: G protein-coupled receptor; glycoprotein; phosphoprotein; transmembrane
protein
F;40-65/Domain: transmembrane #status predicted <TM1>
F;76-97/Domain: transmembrane #status predicted <TM2>
F;112-133/Domain: transmembrane #status predicted <TM3>
F;151-175/Domain: transmembrane #status predicted <TM4>
F;208-226/Domain: transmembrane #status predicted <TM5>
F;243-264/Domain: transmembrane #status predicted <TM6>
F;291-308/Domain: transmembrane #status predicted <TM7>
F;29-276,110-187/Disulfide bonds: #status predicted
F;72,350/Binding site: phosphate (Ser) (covalent) (by casein kinase II) #status predicted
F;145/Binding site: phosphate (Ser) (covalent) (by protein kinase C) #status predicted
F;183,194/Binding site: carbohydrate (Asn) (covalent) #status predicted
F;321/Binding site: phosphate (Thr) (covalent) (by protein kinase C) #status predicted
                           3.4%; Score 12; DB 2; Length 360;
  Best Local Similarity 100.0%; Pred. No. 0.00065;
  Matches 12; Conservative
                                 0; Mismatches
                                                   0; Indels
                                                                  0; Gaps
                                                                              0;
Qу
       72 YLLNLAISDLLF 83
          111111111111
Db
       77 YLLNLAISDLLF 88
```

SEQ ID NO: 6

Result No.	Score	Query Match	Length	DB	ID	Description
1	343	96.6	355	17	AAW03378	CC-chemokine recep
2	343	96.6	355	19	AAW51746	Human C-C chemokin
3	181	51.0	355	17	AAW03376	CC-chemokine recep
4	181	51.0	355	17	AAW03377	CC-chemokine recep
5	181	51.0	355	18	AAW31850	Human eosinophil e
6	181	51.0	355	18	AAW27124	Human chemokine re
7	181	51.0	355	18	AAW10100	Human C-C chemokin
8	181	51.0	355	19	AAW51745	Human C-C chemokin
9	181	51.0	355	22	ABB56341	Non-endogenous hum
10	181	51.0	355	22	AAG80109	Human CCR3 protein
11	181	51.0	356	18	AAW25943	Human CCKR3 chemok
12	134	37.7	355	19	AAW51744	Human C-C chemokin
13	41	11.5	295	22	AAG80106	Human CCR1 protein
14	41	11.5	355	15	AAR52749	C-C chemokine rece
15	41	11.5	355	18	AAW26588	Human MIP-1 alpha/
16	41	11.5	355	18	AAW25751	Human MIP-1alpha/R
17	41	11.5	355	21	AAB20571	Human CC-chemokine
18	34	9.6	34	22	AAG80053	Chemokine peptide
19	31	8.7	31	16	AAR79170	End of third trans

```
RESULT 14
AAR52749
ID
    AAR52749 standard; Protein; 355 AA.
XX
AC
     AAR52749;
XX
DT
     30-JAN-1995 (first entry)
XX
DΕ
     C-C chemokine receptor.
XX
     C-C CKR-1; cytokine; inflammation.
KW
XX
os
     Homo sapiens.
XX
PN
     WO9411504-A.
xx
PD
     26-MAY-1994.
xx
PF
     04-NOV-1993;
                    93WO-US10672.
XX
PR
     10-NOV-1992;
                    92US-0974025.
XX
PΑ
     (GETH ) GENENTECH INC.
\mathbf{x}\mathbf{x}
ΡI
     Horuk R, Neote K, Schall T;
XX
DR
     WPI; 1994-183505/22.
     N-PSDB; AAQ62695.
DR
XX
PT
     New C-C chemokine receptor and nucleic acid - are used to develop
PT
     prods. for use in diagnosis and therapy of inflammation and other
     cytokine-mediated disorders
PT
XX
PS
     Claim 1; Fig 9; 90pp; English.
XX
CC
     The sequence is that of the C-C chemokine receptor. The sequence can
CC
     be used in therapeutic or diagnostic compsns. for inflammation and
CC
     other cytokine mediated disorders.
CC
     See also AAR52750-2.
XX
SQ
     Sequence 355 AA;
```

```
11.5%; Score 41; DB 15; Length 355;
  Ouerv Match
  Best Local Similarity 100.0%; Pred. No. 2.1e-32;
  Matches 41; Conservative
                                0; Mismatches
                                                  0; Indels
                                                                 0; Gaps
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qу
          Db
      115 tglyseiffiilltidrylaivhavfalrartvtfgvitsi 155
RESULT 15
AAW26588
ID
     AAW26588 standard; Protein; 355 AA.
XX
AC
    AAW26588;
XX
     21-JAN-1998 (first entry)
DT
XX
     Human MIP-1 alpha/RANTES receptor.
DE
XX
KW
     Macrophage inflammatory protein-1 alpha; MIP-1 alpha;
     reduced upon activation normal T expressed and secreted; RANTES;
KW
KW
     receptor; cytokine; antiinflammatory; inflammation; human.
XX
os
     Homo sapiens.
XX
ΡN
     US5652133-A.
XX
PD
     29-JUL-1997.
XX
PF
     28-JAN-1993;
                    93US-0012988.
XX
     28-JAN-1993;
                    93US-0012988.
PR
XX
     (USSH ) US DEPT HEALTH & HUMAN SERVICES.
PA
xx
PΙ
     Murphy PM;
XX
DR
     WPI; 1997-392945/36.
DR
     N-PSDB; AAT90384.
XX
PT
     MIP-1-alpha and RANTES receptor nucleic acid - used to develop
PT
     products for the detection of these cytokine(s) and their receptors,
PT
     particularly in inflammatory processes
\mathbf{x}\mathbf{x}
PS
     Claim 2; Column 15-18; 12pp; English.
XX
CC
     This polypeptide comprises a claimed receptor for human macrophage
CC
     inflammatory protein-1 alpha (MIP-1 alpha) and regulated upon
     activation normal T expressed and secreted (RANTES) protein. Also
CC
     claimed are: a nucleic acid (see AAT90384) that encodes the receptor;
CC
CC
     a subsequence of the nucleic acid, having at least 12 contiguous
CC
     nucleotides; a cell transformed or transfected with the nucleic
CC
     acid; and purified MIP-1 alpha/RANTES receptor polypeptide. The
CC
     products can be used for detecting the MIP-1 alpha/RANTES receptor
CC
     and polymorphisms in physiological samples. In addition, the
CC
     receptor can be expressed and used to assay for MIP-la/RANTES in
     biological samples. The quantitation of MIP-1 alpha/RANTES is
CC
     useful for monitoring the levels of these cytokines in a patient.
CC
CC
     Such measurements are useful in following the antiinflammatory
CC
     effects of drugs and prospective usefulness of new antiinflammatory
CC
     agents.
XX
SQ
     Sequence 355 AA;
                         11.5%; Score 41; DB 18; Length 355; 100.0%; Pred. No. 2.1e-32;
  Query Match
  Best Local Similarity
```

Matches 41; Conservative 0; Mismatches 0; Indels

0; Gaps

0;

٩

Result No.	Score	Query Match	Length	DB	ID .	Description
1	181	51.0	355	4	US-08-575-967A-4	Sequence 4, Appli
2	181	51.0	355	4	US-08-847-296B-1	Sequence 1, Appli
3	181	51.0	355	4	US-09-045-583-54	Sequence 54, Appl
4	41	11.5	355	1	US-08-012-988A-2	Sequence 2, Appli
5	41	11.5	355	1	US-08-450-393A-5	Sequence 5, Appli
6	41	11.5	355	4	US-08-446-669-5	Sequence 5, Appli
7	41	11.5	355	4	US-09-045-583-53	Sequence 53, Appl
8	41	11.5	355	4	US-09-239-938-1	Sequence 1, Appli
9	41	11.5	355	5	PCT-US95-00476-5	Sequence 5, Appli
10	31	8.7	31	1	US-08-450-393A-14	Sequence 14, Appl
11	31	8.7	31	4	US-08-446-669-14	Sequence 14, Appl
12	31	8.7	31	5	PCT-US95-00476-14	Sequence 14, Appl
13	22	6.2	31	1	US-08-450-393A-13	Sequence 13, Appl
14	22	6.2	31	4	US-08-446-669-13	Sequence 13, Appl
15	22	6.2	31	5	PCT-US95-00476-13	Sequence 13, Appl
16	22	6.2	344	3	US-08-466-343D-9	Sequence 9, Appli
17	22	6.2	347	1	US-08-461-244-3	Sequence 3, Appli
18	22	6.2	352	3	US-08-466-343D-2	Sequence 2, Appli
19	22	6.2	352	4	US-09-045-583-52	Sequence 52, Appl
20	22	6.2	360	1	US-08-450-393A-4	Sequence 4, Appli

```
RESULT 4
US-08-012-988A-2
; Sequence 2, Application US/08012988A
; Patent No. 5652133
; GENERAL INFORMATION:
    APPLICANT: Murphy, Philip M.
TITLE OF INVENTION: Cloning and Expression of Human
    TITLE OF INVENTION: Macrophage Inflammatory Protein-1 alpha (MIP-1
    TITLE OF INVENTION: alpha)/RANTES Receptor
    NUMBER OF SEQUENCES: 2
    CORRESPONDENCE ADDRESS:
      ADDRESSEE: Townsend and Townsend Khourie and Crew
      STREET: One Market Plaza, Steuart Tower, Suite 2000
      CITY: San Francisco
      STATE: California
      COUNTRY: USA
      ZIP: 94610
    COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
      OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
      APPLICATION NUMBER: US/08/012,988A
      FILING DATE: 19930128
      CLASSIFICATION: 435
    ATTORNEY/AGENT INFORMATION:
      NAME: Weber,, Kenneth A.
      REGISTRATION NUMBER: 31,677
      REFERENCE/DOCKET NUMBER: 15280-118
    TELECOMMUNICATION INFORMATION:
      TELEPHONE: 415-543-9600
      TELEFAX: 415-543-5043
  INFORMATION FOR SEQ ID NO: 2:
    SEQUENCE CHARACTERISTICS:
      LENGTH: 355 amino acids
      TYPE: AMINO ACID
```

```
TOPOLOGY: linear
    MOLECULE TYPE: protein
US-08-012-988A-2
                         11.5%; Score 41; DB 1; Length 355;
  Best Local Similarity 100.0%; Pred. No. 3.7e-31;
  Matches 41; Conservative 0; Mismatches
                                                0; Indels
                                                                0; Gaps
                                                                            0;
Qу
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
          Db
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
RESULT 5
US-08-450-393A-5
; Sequence 5, Application US/08450393A
; Patent No. 5707815
; GENERAL INFORMATION:
    APPLICANT: Charo, Israel
APPLICANT: Coughlin, Shaun
    TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT
    TITLE OF INVENTION: PROTEIN RECEPTORS NUMBER OF SEQUENCES: 14
    CORRESPONDENCE ADDRESS:
      ADDRESSEE: Cooley Godward Castro Huddleson & Tatum
      STREET: 5 Palo Alto Square
      CITY: Palo Alto
STATE: California
      COUNTRY: USA
      ZIP: 94306-2155
    COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
      APPLICATION NUMBER: US/08/450,393A FILING DATE: May 25, 1995
      CLASSIFICATION: 424
    ATTORNEY/AGENT INFORMATION:
      NAME: Cserr, Luann
      REGISTRATION NUMBER: 31,822
      REFERENCE/DOCKET NUMBER: UCAL-237/02US
    TELECOMMUNICATION INFORMATION:
      TELEPHONE: 415-843-5165
      TELEFAX: 415-8857-0663
      TELEX: 380816CooleyPA
  INFORMATION FOR SEQ ID NO: 5:
    SEQUENCE CHARACTERISTICS:
      LENGTH: 355 amino acids
      TYPE: amino acid
      TOPOLOGY: linear
    MOLECULE TYPE: protein
    HYPOTHETICAL: NO
    ANTI-SENSE: NO
US-08-450-393A-5
                         11.5%; Score 41; DB 1; Length 355;
  Best Local Similarity 100.0%; Pred. No. 3.7e-31;
 Matches 41; Conservative
                               0; Mismatches
                                                0; Indels
                                                                0; Gaps
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qу
```

115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155

Db

```
RESULT 12
PCT-US95-00476-14
; Sequence 14, Application PC/TUS9500476
: GENERAL INFORMATION:
    APPLICANT: The Regents of the University of California
     TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT
    TITLE OF INVENTION: PROTEIN RECEPTORS NUMBER OF SEQUENCES: 14
    CORRESPONDENCE ADDRESS:
       ADDRESSEE: Robbins, Berliner & Carson
       STREET: 201 N. Figueroa Street, 5th Floor
       CITY: Los Angeles
STATE: California
       COUNTRY: USA
       ZIP: 90012-2628
    COMPUTER READABLE FORM:
       MEDIUM TYPE: Floppy disk
       COMPUTER: IBM PC compatible
       OPERATING SYSTEM: PC-DOS/MS-DOS
       SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
      APPLICATION NUMBER: PCT/US95/00476
       FILING DATE:
      CLASSIFICATION:
    ATTORNEY/AGENT INFORMATION:
     NAME: Berliner, Robert
       REGISTRATION NUMBER: 20,121
       REFERENCE/DOCKET NUMBER: 5555-291
    TELECOMMUNICATION INFORMATION:
       TELEPHONE: 310-977-1001
       TELEFAX: 310-977-1003
       TELEX:
  INFORMATION FOR SEQ ID NO: 14:
    SEQUENCE CHARACTERISTICS:
      LENGTH: 31 amino acids
       TYPE: amino acid
       TOPOLOGY: linear
    MOLECULE TYPE: peptide
PCT-US95-00476-14
 Query Match 8.7%; Score 31; DB 5; Length 31; Best Local Similarity 100.0%; Pred. No. 1.3e-22;
 Matches 31; Conservative
                                0; Mismatches
                                                   0; Indels
                                                                  0; Gaps
                                                                              0;
      121 IFFIILLTIDRYLAIVHAVFALRARTVTFGV 151
Qу
          Db
        1 IFFIILLTIDRYLAIVHAVFALRARTVTFGV 31
```

		*				
Result		Query				
No.	Score	Match	Length	DB	ID	Description
				- 		
1	181	51.0	355	1	CKR3_HUMAN	P51677 homo sapien
2	64	18.0	355	1	CKR3_MACMU	P56483 macaca mula
3	55	15.5	355	1	CKR3_CERAE	P56492 cercopithec
4	41	11.5	355	1	CKR1_HUMAN	P32246 homo sapien
5	41	11.5	355	1	CKR1_MACMU	P56482 macaca mula
6	33	9.3	355	1	CKR1_MOUSE	P51675 mus musculu
7	33	9.3	359	1	CKR3_MOUSE	P51678 mus musculu
8	29	8.2	358	1	CKR3_CAVPO	Q9z2i3 cavia porce
9	23	6.5	359	1	CKR3_RAT	O54814 rattus norv
10	22	6.2	352	1	CKR5_CERAE	P56493 cercopithec
11	22	6.2	352	1	CKR5_CERTO	O62743 cercocebus
12	22	6.2	352	1	CKR5_GORGO	P56439 gorilla gor
13	22	6.2	352	1	CKR5_HYLLE	O97883 hylobates l
14	22	6.2	352	1	CKR5_MACMU	P79436 macaca mula

```
RESULT
CKR1_HUMAN
    CKR1_HUMAN
                   STANDARD;
                                  PRT;
                                         355 AA.
ID
AC
    P32246:
    01-OCT-1993 (Rel. 27, Created)
    01-OCT-1993 (Rel. 27, Last sequence update)
DT
    16-OCT-2001 (Rel. 40, Last annotation update)
DT
    C-C chemokine receptor type 1 (C-C CKR-1) (CC-CKR-1) (CCR-1) (CCR1)
DE
DE
     (Macrophage inflammatory protein-1 alpha receptor) (MIP-lalpha-R)
DΕ
     (RANTES-R) (HM145) (LD78 receptor).
GN
    CCR1 OR CMKBR1 OR CMKR1.
os
    Homo sapiens (Human).
    Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC
    Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
OC
ox
    NCBI_TaxID=9606;
RN
     [1]
RP
    SEQUENCE FROM N.A.
RX
    MEDLINE=93161416; PubMed=7679328;
RA
    Neote K., Digregorio D., Mak J.Y., Horuk R., Schall T.J.;
    "Molecular cloning, functional expression, and signaling
RT
RT
    characteristics of a C-C chemokine receptor.";
    Cell 72:415-425(1993).
RL
RN
    SEQUENCE FROM N.A.
RР
RX
    MEDLINE=93240122; PubMed=7683036;
    Gao J.-L., Kuhns D., Tiffany H.L., McDermott D., Li X., Francke U.,
RA
RA
    Murphy P.M.;
RT
     "Structure and functional expression of the human macrophage
    inflammatory protein 1 alpha/RANTES receptor.";
RT
    J. Exp. Med. 177:1421-1427(1993).
RL
RN
    [3]
    SEQUENCE FROM N.A.
RÞ
RC
    TISSUE=Monocytes;
    MEDLINE=94092629; PubMed=7505609;
RX
RA
    Nomura H., Nielsen B.W., Matsushima K.;
    "Molecular cloning of cDNAs encoding a LD78 receptor and putative
RT
RT
    leukocyte chemotactic peptide receptors.";
RL
    Int. Immunol. 5:1239-1249(1993).
     -!- FUNCTION: RECEPTOR FOR A C-C TYPE CHEMOKINE. BINDS TO MIP-1-ALPHA,
CC
        MIP-1 DELTA, RANTES, AND MCP-3 AND, LESS EFFICIENTLY, TO MIP-1-
CC
CC
        BETA OR MCP-1 AND SUBSEQUENTLY TRANSDUCES A SIGNAL BY INCREASING
CC
        THE INTRACELLULAR CALCIUM IONS LEVEL. RESPONSIBLE FOR AFFECTING
CC
        STEM CELL PROLIFERATION.
CC
    -!- SUBCELLULAR LOCATION: Integral membrane protein.
    -!- TISSUE SPECIFICITY: WIDELY EXPRESSED IN DIFFERENT HEMATOPOIETIC
CC
        CELLS.
CC
    -!- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.
CC
CC
                    -----
    This SWISS-PROT entry is copyright. It is produced through a collaboration
CC
    between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC
    the European Bioinformatics Institute. There are no restrictions on its
CC
CC
    use by non-profit institutions as long as its content is in no way
CC
    modified and this statement is not removed. Usage by and for commercial
    entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC
CC
    or send an email to license@isb-sib.ch).
                               _____
CC
    EMBL; L09230; AAA58408.1; -.
DR
    EMBL; L10918; AAA36543.1; -.
DR
    EMBL; D10925; BAA01723.1; -.
DR
    PIR; A45177; A45177.
DR
DR
    GCRDb; GCR_0498; -.
DR
    GCRDb; GCR 0557; -.
DR
    GCRDb; GCR_0573; -.
DR
    MIM; 601159; -.
    InterPro; IPR000276; GPCR Rhodpsn.
DR
    Pfam; PF00001; 7tm_1; 1.
DR
    PRINTS; PR00237; GPCRRHODOPSN.
DR
    PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR
DR
    PROSITE; PS50262; G PROTEIN RECEP F1 2; 1.
    G-protein coupled receptor; Transmembrane; Glycoprotein.
```

```
FT
     DOMAIN
                         34
                                   EXTRACELLULAR (POTENTIAL).
FT
     TRANSMEM
                  35
                         60
                                   1 (POTENTIAL).
FT
     DOMAIN
                  61
                         64
                                   CYTOPLASMIC (POTENTIAL).
FT
     TRANSMEM
                  65
                         91
                                   2 (POTENTIAL).
                                   EXTRACELLULAR (POTENTIAL).
                  92
FT
     DOMATN
                        107
     TRANSMEM
                 108
                        129
                                   3 (POTENTIAL).
FT
                                  CYTOPLASMIC (POTENTIAL).
     DOMAIN
                 130
                        146
FT
FT
     TRANSMEM
                 147
                        171
                                   4 (POTENTIAL).
     DOMAIN
                 172
                                  EXTRACELLULAR (POTENTIAL).
FT
                        197
FT
     TRANSMEM
                 198
                        223
                                   5 (POTENTIAL).
FT
     DOMAIN
                 224
                        239
                                  CYTOPLASMIC (POTENTIAL).
     TRANSMEM
FT
                 240
                        264
                                  6 (POTENTIAL).
                                  EXTRACELLULAR (POTENTIAL).
FT
     DOMAIN
                 265
                        281
FΤ
     TRANSMEM
                 282
                        305
                                  7 (POTENTIAL).
FT
     DOMAIN
                 306
                        355
                                  CYTOPLASMIC (POTENTIAL).
FT
     CARBOHYD
                  5
                                  \mbox{N-LINKED} (GLCNAC. . .) (POTENTIAL).
                         5
FT
     DISULFID
                 106
                        183
                                  BY SIMILARITY.
                                  E \rightarrow D (IN REF. 3).
FT
     CONFLICT
                 337
                        337
                355 AA; 41172 MW; B2C100FFED275985 CRC64;
SO
     SEQUENCE
  Query Match
                          11.5%; Score 41; DB 1; Length 355;
  Best Local Similarity 100.0%; Pred. No. 4.2e-31;
           41; Conservative
                                 0; Mismatches
                                                    0: Indels
                                                                  0; Gaps
Qу
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
          Db
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
RESULT
CKR3_MOUSE
ID
     CKR3 MOUSE
                    STANDARD;
                                    PRT:
                                          359 AA.
AC
     P51678;
     01-OCT-1996 (Rel. 34, Created)
DT
     01-OCT-1996 (Rel. 34, Last sequence update)
DT
DT
     30-MAY-2000 (Rel. 39, Last annotation update)
DE
     Probable C-C chemokine receptor type 3 (C-C CKR-3) (CC-CKR-3) (CCR-3)
     (CCR3) (CKR3) (Macrophage inflammatory protein-1 alpha receptor-like
DE
     2) (MIP-1 alpha RL2).
DE
GN
     CCR3 OR CMKBR3 OR CMKBR1L2.
os
     Mus musculus (Mouse).
OC
     Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC
     Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.
ox
     NCBI TaxID=10090;
RN
     [1]
RP
     SEQUENCE FROM N.A.
RC
     STRAIN=129/SV;
RX
     MEDLINE=96072806; PubMed=7594543;
RA
     Post T.W., Bozic C.R., Rothenberg M.E., Luster A.D., Gerard N.,
RA
RТ
     "Molecular characterization of two murine eosinophil beta chemokine
RT
     receptors.";
RL
     J. Immunol. 155:5299-5305(1995).
RN
     [2]
RP
     SEQUENCE FROM N.A.
RC
     STRAIN=129/SVJ;
RX
     MEDLINE=95340546; PubMed=7542241;
RA
     Gao J.-L., Murphy P.M.;
RT
     "Cloning and differential tissue-specific expression of three mouse
RT
     beta chemokine receptor-like genes, including the gene for a
     functional macrophage inflammatory protein-1 alpha receptor.";
RT
RL
     J. Biol. Chem. 270:17494-17501(1995).
CC
     -!- FUNCTION: RECEPTOR FOR A C-C TYPE CHEMOKINE. BINDS TO EOTAXIN,
         MCP-3, MCP-4 AND RANTES AND SUBSEQUENTLY TRANSDUCES A SIGNAL BY
CC
CC
         INCREASING THE INTRACELLULAR CALCIUM IONS LEVEL.
CC
     -!- SUBCELLULAR LOCATION: Integral membrane protein.
    -!- TISSUE SPECIFICITY: DETECTED IN SKELETAL MUSCLE AND IN TRACE
CC
CC
         AMOUNTS IN LEUKOCYTES.
     -!- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.
```

```
This SWISS-PROT entry is copyright. It is produced through a collaboration
CC
    between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC
CC
    the European Bioinformatics Institute. There are no restrictions on its
    use by non-profit institutions as long as its content is in no way
CC
CC
    modified and this statement is not removed. Usage by and for commercial
    entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC
CC
    or send an email to license@isb-sib.ch).
CC
DR
    EMBL; U29677; AAA86118.1; -.
DR
    EMBL; U28406; AAA89155.1; -.
    GCRDb; GCR_1673; -.
DR
    GCRDb; GCR_1695; -.
DR
    MGD; MGI:104616; Cmkbr112.
DR
DR
    InterPro; IPR000276; GPCR_Rhodpsn.
    Pfam; PF00001; 7tm_1; 1.
DR
DR
    PRINTS; PR00237; GPCRRHODOPSN.
DR
    PROSITE; PS00237; G PROTEIN RECEP F1 1; 1.
    PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
DR
    G-protein coupled receptor; Transmembrane.
KW
                                EXTRACELLULAR (POTENTIAL).
    DOMAIN
FT
                       38
                 1
FT
    TRANSMEM
                 39
                        64
                                1 (POTENTIAL).
    DOMAIN
                                CYTOPLASMIC (POTENTIAL).
FT
                 65
                       68
FT
    TRANSMEM
                 69
                       95
                                2 (POTENTIAL).
FT
    DOMAIN
                96
                       111
                                EXTRACELLULAR (POTENTIAL).
FT
    TRANSMEM
                112
                       133
                                3 (POTENTIAL).
FT
    DOMAIN
                134
                      150
                                CYTOPLASMIC (POTENTIAL).
    TRANSMEM
                                4 (POTENTIAL).
FT
                151
                      175
    DOMAIN
FT
                176
                      201
                                EXTRACELLULAR (POTENTIAL).
    TRANSMEM
FТ
                                5 (POTENTIAL).
                202
                      227
    DOMAIN
                228
                                CYTOPLASMIC (POTENTIAL).
FT
                       243
FT
    TRANSMEM
                                6 (POTENTIAL).
                244
                      268
FT
    DOMAIN
                269
                       285
                                EXTRACELLULAR (POTENTIAL).
FT
    TRANSMEM
                286
                      309
                                7 (POTENTIAL).
                                CYTOPLASMIC (POTENTIAL).
FT
    DOMAIN
                310
                      359
                       187
FT
    DISULFID
                110
                                BY SIMILARITY.
    CONFLICT
                270
                      270
                                R \rightarrow S (IN REF. 2).
FT
    SEQUENCE
              359 AA; 41825 MW; AC11ED66E283CEAF CRC64;
 Query Match
                         9.3%; Score 33; DB 1; Length 359;
 Best Local Similarity 100.0%; Pred. No. 1.6e-23;
 Matches 33; Conservative 0; Mismatches
                                                0: Indels
                                                               0: Gaps
                                                                           0:
Qу
     117 LYSEIFFIILLTIDRYLAIVHAVFALRARTVTF 149
         Db
     121 LYSEIFFIILLTIDRYLAIVHAVFALRARTVTF 153
```

CC

	8				
	Query				
Score	Match	Length	DB	ID	Description
72	20.3	72	4	Q96T96	O96t96 homo sapien
56	15.8	56	4	Q96T97	Q96t97 homo sapien
51	14.4	355	6	Q9BDS8	Q9bds8 macaca fasc
41	11.5	355	6	Q9MYJ8	Q9myj8 callithrix
35	9.9	355	11	Q91VP9	Q91vp9 mus musculu
35	9.9	358	6	Q9N0M0	Q9n0m0 ovis aries
26	7.3	351	6	Q9MYJ9	Q9myj9 oryctolagus
25	7.0	355	11	Q9JLY8	Q9jly8 rattus norv
22	6.2	316	6	Q9TUV7	Q9tuv7 saguinus sp
22	6.2	334	6	Q9TUQ7	Q9tuq7 erythrocebu
22	6.2	339	6	Q9TQX3	Q9tqx3 mandrillus
22	6.2	339	6	Q9TQX2	Q9tqx2 erythrocebu
22	6.2	339	6	Q9TQW4	Q9tqw4 pan troglod
22	6.2	339	6	Q9TQW2	Q9tqw2 pongo pygma
22	6.2	339	6	Q9TQW0	Q9tqw0 hylobates c
	56 51 41 35 35 26 25 22 22 22 22 22 22	Query Score Match 72 20.3 56 15.8 51 14.4 41 11.5 35 9.9 35 9.9 26 7.3 25 7.0 22 6.2 22 6.2 22 6.2 22 6.2 22 6.2 22 6.2 22 6.2	Query Score Match Length 72 20.3 72 56 15.8 56 51 14.4 355 41 11.5 355 35 9.9 358 26 7.3 351 25 7.0 355 22 6.2 336 22 6.2 339 22 6.2 339 22 6.2 339 22 6.2 339	Query Score Match Length DB 72 20.3 72 4 56 15.8 56 4 51 14.4 355 6 41 11.5 355 6 35 9.9 355 11 35 9.9 358 6 26 7.3 351 6 25 7.0 355 11 22 6.2 316 6 22 6.2 334 6 22 6.2 339 6 22 6.2 339 6 22 6.2 339 6 22 6.2 339 6	Query Score Match Length DB ID 72 20.3 72 4 Q96T96 56 15.8 56 4 Q96T97 51 14.4 355 6 Q9BDS8 41 11.5 355 6 Q9MYJ8 35 9.9 355 11 Q91VP9 35 9.9 358 6 Q9NOMO 26 7.3 351 6 Q9MYJ9 25 7.0 355 11 Q9JLY8 22 6.2 316 6 Q9TUV7 22 6.2 334 6 Q9TUQ7 22 6.2 339 6 Q9TQX2 22 6.2 339 6 Q9TQW4 22 6.2 339 6 Q9TQW4

16	22	6.2	339	6	Q9TQV6	Q9tqv6 colobus gue
17	22	6.2	339	6	Q9TQV5	Q9tqv5 saguinus sp
18	22	6.2	339	6	Q9TQV3	Q9tqv3 cercopithec

```
      16
      22
      6.2
      339
      6
      Q9TQV6
      Q9tqv6 colobus gue

      17
      22
      6.2
      339
      6
      Q9TQV5
      Q9tqv5 saguinus sp

      18
      22
      6.2
      339
      6
      Q9TQV3
      Q9tqv3 cercopithec
```

SEQ ID NO: 4

		*				
Result		Query				
No.	Score	Match	Length	ĎВ	ID	Description
	255	100.0				OC shareline we see
1	355	100.0	355	17	AAW03377	CC-chemokine recep
2	355	100.0	355	18	AAW31850	Human eosinophil e
3	355	100.0	355	18	AAW27124	Human chemokine re
4	355	100.0	355	19	AAW51745	Human C-C chemokin
5	355	100.0	355	22	AAG80109	Human CCR3 protein
6	281	79.2	356	18	AAW25943	Human CCKR3 chemok
7	275	77.5	355	17	AAW03376	CC-chemokine recep
8	275	77.5	355	18	AAW10100	Human C-C chemokin
9	254	71.5	355	22	ABB56341	Non-endogenous hum
10	228	64.2	355	19	AAW51744	Human C-C chemokin
11	181	51.0	355	17	AAW03378	CC-chemokine recep
12	181	51.0	355	19	AAW51746	Human C-C chemokin
13	41	11.5	295	22	AAG80106	Human CCR1 protein
14	41	11.5	355	15	AAR52749	C-C chemokine rece
15	41	11.5	355	18	AAW26588	Human MIP-1 alpha/
16	41	11.5	355	18	AAW25751	Human MIP-lalpha/R
17	41	11.5	355	21	AAB20571	Human CC-chemokine
18	34	9.6	34	22	AAG80053	Chemokine peptide
19	32	9.0	32	22	AAG80082	Chemokine CCR3 ext
20	31	8.7	31	16	AAR79170	End of third trans
21	27	7.6	28	20	AAY39255	G-protein coupled
22	26	7.3	28	20	AAY39256	G-protein coupled

```
RESULT 14
AAR52749
ID
    AAR52749 standard; Protein; 355 AA.
XX
АC
    AAR52749;
XX
DT
     30-JAN-1995 (first entry)
XX
DE
     C-C chemokine receptor.
xx
KW
     C-C CKR-1; cytokine; inflammation.
xx
os
     Homo sapiens.
XX
PN
     WO9411504-A.
XX
PD
     26-MAY-1994.
XX
PF
     04-NOV-1993;
                    93WO-US10672.
XX
PR
     10-NOV-1992;
                    92US-0974025.
XX
PA
     (GETH ) GENENTECH INC.
\mathbf{x}\mathbf{x}
PΙ
    Horuk R, Neote K, Schall T;
XX
DR
     WPI; 1994-183505/22.
DR
     N-PSDB; AAQ62695.
XX
PT
     New C-C chemokine receptor and nucleic acid - are used to develop
PΤ
     prods. for use in diagnosis and therapy of inflammation and other
PT
     cytokine-mediated disorders
XX
PS
     Claim 1; Fig 9; 90pp; English.
XX
```

```
CC
     The sequence is that of the C-C chemokine receptor. The sequence can
     be used in therapeutic or diagnostic compsns. for inflammation and
CC
CC
     other cytokine mediated disorders.
CC
     See also AAR52750-2.
XX
SQ
     Sequence
                355 AA;
  Query Match
                          11.5%; Score 41; DB 15; Length 355;
  Best Local Similarity 100.0%; Pred. No. 1.2e-31;
                                                   0; Indels
           41; Conservative
                                0; Mismatches
                                                                 0; Gaps
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qy
          Db
      115 tglyseiffiilltidrylaivhavfalrartvtfgvitsi 155
RESULT 15
AAW26588
     AAW26588 standard; Protein; 355 AA.
ID
XX
AC
     AAW26588:
XX
DT
     21-JAN-1998 (first entry)
XX
DE
     Human MIP-1 alpha/RANTES receptor.
XX
KW
     Macrophage inflammatory protein-1 alpha; MIP-1 alpha;
KW
     reduced upon activation normal T expressed and secreted; RANTES;
KW
     receptor; cytokine; antiinflammatory; inflammation; human.
XX
os
     Homo sapiens.
\mathbf{x}\mathbf{x}
PN
     US5652133-A.
XX
PΠ
     29-JUL-1997.
XX
PF
     28-JAN-1993;
                    93US-0012988.
XX
PR
     28-JAN-1993:
                    93US-0012988.
XX
PA
     (USSH ) US DEPT HEALTH & HUMAN SERVICES.
XX
ΡI
     Murphy PM;
XX
DR
     WPI; 1997-392945/36.
     N-PSDB; AAT90384.
DR
XX
PT
     MIP-1-alpha and RANTES receptor nucleic acid - used to develop
PT
     products for the detection of these cytokine(s) and their receptors,
PT
     particularly in inflammatory processes
XX
PS
     Claim 2; Column 15-18; 12pp; English.
XX
CC
     This polypeptide comprises a claimed receptor for human macrophage
CC
     inflammatory protein-1 alpha (MIP-1 alpha) and regulated upon
CC
     activation normal T expressed and secreted (RANTES) protein. Also
CC
     claimed are: a nucleic acid (see AAT90384) that encodes the receptor;
CC
     a subsequence of the nucleic acid, having at least 12 contiguous
CC
     nucleotides; a cell transformed or transfected with the nucleic
CC
     acid; and purified MIP-1 alpha/RANTES receptor polypeptide. The
CC
     products can be used for detecting the MIP-1 alpha/RANTES receptor
CC
     and polymorphisms in physiological samples. In addition, the
CC
     receptor can be expressed and used to assay for MIP-la/RANTES in
CC
     biological samples. The quantitation of MIP-1 alpha/RANTES is
CC
     useful for monitoring the levels of these cytokines in a patient.
CC
     Such measurements are useful in following the antiinflammatory
CC
     effects of drugs and prospective usefulness of new antiinflammatory
CC
```

```
11.5%; Score 41; DB 18; Length 355;
  Ouery Match
  Best Local Similarity 100.0%; Pred. No. 1.2e-31;
                               0; Mismatches
                                                     Indels
                                                                0; Gaps
          41; Conservative
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qу
          Db
      115 tglyseiffiilltidrylaivhavfalrartvtfgvitsi 155
                                   SUMMARIES
                윢
Result
               Query
        Score Match Length DB ID
                                                          Description
  No.
                        355 4 US-08-575-967A-4
                                                           Sequence 4, Appli
          355 100.0
    1
    2
          355
               100.0
                        355 4 US-08-847-296B-1
                                                           Sequence 1, Appli
                        355 4 US-09-045-583-54
          355 100.0
                                                           Sequence 54, Appl
    3
    4
           41
               11.5
                        355 1 US-08-012-988A-2
                                                           Sequence 2, Appli
                        355 1 US-08-450-393A-5
355 4 US-08-446-669-5
    5
           41
                11.5
                                                           Sequence 5, Appli
    6
           41
                11.5
                                                           Sequence 5, Appli
    7
           41
                11.5
                        355 4 US-09-045-583-53
                                                           Sequence 53, Appl
                        355 4 US-09-239-938-1
                                                           Sequence 1, Appli
    8
                11.5
           41
    9
           41
                11.5
                        355 5 PCT-US95-00476-5
                                                           Sequence 5, Appli
                         31 1 US-08-450-393A-14
   10
                 8.7
                                                           Sequence 14, Appl
           31
    11
           31
                 8.7
                         31 4 US-08-446-669-14
                                                           Sequence 14, Appl
                         31 5 PCT-US95-00476-14
31 1 US-08-450-393A-13
    12
                                                           Sequence 14, Appl
           31
                 8.7
   13
           22
                 6.2
                                                           Sequence 13, Appl
    14
           22
                 6.2
                         31 4 US-08-446-669-13
                                                           Sequence 13, Appl
   15
           22
                 6.2
                         31 5 PCT-US95-00476-13
                                                           Sequence 13, Appl
   16
           22
                 6.2
                        344
                            3
                                US-08-466-343D-9
                                                           Sequence 9, Appli
                        347 1 US-08-461-244-3
   17
           22
                 6.2
                                                           Sequence 3, Appli
                        352 3 US-08-466-343D-2
                                                           Sequence 2, Appli
   18
                 6.2
RESULT
US-08-012-988A-2
; Sequence 2, Application US/08012988A
; Patent No. 5652133
  GENERAL INFORMATION:
    APPLICANT: Murphy, Philip M.
    TITLE OF INVENTION: Cloning and Expression of Human
    TITLE OF INVENTION: Macrophage Inflammatory Protein-1 alpha (MIP-1
    TITLE OF INVENTION: alpha) / RANTES Receptor
    NUMBER OF SEQUENCES: 2
    CORRESPONDENCE ADDRESS:
      ADDRESSEE: Townsend and Townsend Khourie and Crew
      STREET: One Market Plaza, Steuart Tower, Suite 2000
      CITY: San Francisco
      STATE: California
      COUNTRY: USA
      ZIP: 94610
    COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
      OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
    CURRENT APPLICATION DATA:
      APPLICATION NUMBER: US/08/012,988A
      FILING DATE: 19930128
      CLASSIFICATION: 435
    ATTORNEY/AGENT INFORMATION:
      NAME: Weber,, Kenneth A.
      REGISTRATION NUMBER: 31,677
      REFERENCE/DOCKET NUMBER: 15280-118
    TELECOMMUNICATION INFORMATION:
```

Sequence 355 AA;

TELEPHONE: 415-543-9600 TELEFAX: 415-543-5043

```
INFORMATION FOR SEQ ID NO: 2:
     SEQUENCE CHARACTERISTICS:
      LENGTH: 355 amino acids
       TYPE: AMINO ACID
      TOPOLOGY: linear
     MOLECULE TYPE: protein
US-08-012-988A-2
  Query Match 11.5%; Score 41; DB 1; Length 355; Best Local Similarity 100.0%; Pred. No. 6.7e-31;
  Matches 41; Conservative 0; Mismatches 0; Indels
                                                                 0; Gaps
                                                                             0;
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qy
          115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Db
RESULT 5
US-08-450-393A-5
; Sequence 5, Application US/08450393A
; Patent No. 5707815
; GENERAL INFORMATION:
    APPLICANT: Charo, Israel APPLICANT: Coughlin, Shaun
    TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT
    TITLE OF INVENTION: PROTEIN RECEPTORS NUMBER OF SEQUENCES: 14
     CORRESPONDENCE ADDRESS:
      ADDRESSEE: Cooley Godward Castro Huddleson & Tatum
      STREET: 5 Palo Alto Square
      CITY: Palo Alto
STATE: California
      COUNTRY: USA
      ZIP: 94306-2155
     COMPUTER READABLE FORM:
      MEDIUM TYPE: Floppy disk
      COMPUTER: IBM PC compatible
      OPERATING SYSTEM: PC-DOS/MS-DOS
      SOFTWARE: PatentIn Release #1.0, Version #1.25
     CURRENT APPLICATION DATA:
      APPLICATION NUMBER: US/08/450,393A
      FILING DATE: May 25, 1995
      CLASSIFICATION: 424
    ATTORNEY/AGENT INFORMATION:
      NAME: Cserr, Luann
      REGISTRATION NUMBER: 31,822
      REFERENCE/DOCKET NUMBER: UCAL-237/02US
    TELECOMMUNICATION INFORMATION:
      TELEPHONE: 415-843-5165
      TELEFAX: 415-8857-0663
      TELEX: 380816CooleyPA
  INFORMATION FOR SEQ ID NO: 5:
    SEQUENCE CHARACTERISTICS:
      LENGTH: 355 amino acids
      TYPE: amino acid
      TOPOLOGY: linear
    MOLECULE TYPE: protein
    HYPOTHETICAL: NO
    ANTI-SENSE: NO
US-08-450-393A-5
 Query Match 11.5%; Score 41; DB 1; Length 355; Best Local Similarity 100.0%; Pred. No. 6.7e-31;
 Matches 41; Conservative
                              0; Mismatches 0; Indels
                                                                 0; Gaps
Qy
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
          Db
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
```

```
PCT-US95-00476-5
; Sequence 5, Application PC/TUS9500476
   GENERAL INFORMATION:
     APPLICANT: The Regents of the University of California
    TITLE OF INVENTION: MAMMALIAN MONOCYTE CHEMOATTRACTANT TITLE OF INVENTION: PROTEIN RECEPTORS
NUMBER OF SEQUENCES: 14
     CORRESPONDENCE ADDRESS:
       ADDRESSEE: Robbins, Berliner & Carson
       STREET: 201 N. Figueroa Street, 5th Floor
      CITY: Los Angeles
       STATE: California
       COUNTRY: USA
       ZIP: 90012-2628
     COMPUTER READABLE FORM:
       MEDIUM TYPE: Floppy disk
       COMPUTER: IBM PC compatible
OPERATING SYSTEM: PC-DOS/MS-DOS
       SOFTWARE: PatentIn Release #1.0, Version #1.25
     CURRENT APPLICATION DATA:
      APPLICATION NUMBER: PCT/US95/00476
      FILING DATE:
       CLASSIFICATION:
     ATTORNEY/AGENT INFORMATION:
      NAME: Berliner, Robert
       REGISTRATION NUMBER: 20,121
      REFERENCE/DOCKET NUMBER: 5555-291
     TELECOMMUNICATION INFORMATION:
      TELEPHONE: 310-977-1001
       TELEFAX: 310-977-1003
      TELEX:
   INFORMATION FOR SEQ ID NO: 5:
     SEQUENCE CHARACTERISTICS:
      LENGTH: 355 amino acids
      TYPE: amino acid
      TOPOLOGY: linear
     MOLECULE TYPE: protein
     HYPOTHETICAL: NO
     ANTI-SENSE: NO
PCT-US95-00476-5
                          11.5%; Score 41; DB 5; Length 355;
  Query Match
  Best Local Similarity 100.0%; Pred. No. 6.7e-31;
  Matches 41; Conservative 0; Mismatches
                                                 0; Indels
                                                                  0; Gaps
                                                                              0;
Qу
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
          Db
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
                                    SUMMARIES
                 ٩.
R
```

RESULT 9

		ס				
Result		Query				
No.	Score	Match	Length	DB	ID	Description
1	275	77.5	355	2	G02436	chemokine (C-C) re
2	41	11.5	355	2	A45177	chemokine (C-C) re
3	33	9.3	355	2	149339	macrophage inflamm
4	33	9.3	359	2	I49341	MIP-1 alpha recept
5	22	6.2	360	2	JC2443	chemokine (C-C) re
6	22	6.2	374	2	I38450	chemokine (C-C) re
7	16	4.5	383	2	S55594	G protein-coupled
8	14	3.9	352	2	A43113	chemokine (C-C) re
9	12	3.4	356	2	I49340	MIP-1 alpha recept
10	12	3.4	360	2	A57160	chemokine (C-C) re

```
360 2 JC4587
                                                       chemokine (C-C) re
11
       12
             3.4
                     308 2 150241
12
       10
             2.8
                                                       G protein-coupled
                     327 2 S56162
13
       10
             2.8
                                                       MDCR15 protein - h
                                                       G protein-coupled
                     355 2
                             JC5067
14
       10
              2.8
                     372 2 S26667
                                                       G protein-coupled
15
       10
             2.8
       10
             2.8
                     374 2 S42628
                                                       G protein-coupled
16
                    374 2 S32785
177 2 PH1372
17
       10
                                                       G protein-coupled
             2.8
18
        9
             2.5
                                                       angiotensin II rec
19
             2.5
                     350 2 A39445
                                                       interleukin-8 rece
20
             2.5
                     352 2 G00048
                                                       fusin (LESTRA) - c
```

```
RESULT
G02436
chemokine (C-C) receptor 3 - human
N; Alternate names: C-C CKR-3
C; Species: Homo sapiens (man)
C;Date: 21-Dec-1996 #sequence_revision 06-Jun-1997 #text_change 04-Mar-2000
C; Accession: G02436; A57237
R:Ponath, P.D.
submitted to the EMBL Data Library, February 1996
A; Reference number: H01272
A; Accession: G02436
A; Status: translated from GB/EMBL/DDBJ
A; Molecule type: DNA
A; Residues: 1-355 < PON>
A;Cross-references: EMBL:U49727; NID:g1477560; PIDN:AAB09726.1; PID:g1477561
R; Combadiere, C.; Ahuja, S.K.; Murphy, P.M.
J. Biol. Chem. 270, 16491-16494, 1995
A; Title: Cloning and functional expression of a human eosinophil CC chemokine receptor.
A; Reference number: A57237; MUID: 95348056
A; Accession: A57237
A; Status: nucleic acid sequence not shown
A; Molecule type: mRNA
A; Residues: 1-106, 'N', 108-275, 'S', 277-280, 'R', 282-355 < COM>
A; Cross-references: GB:U28694; NID:g1199579; PIDN:AAC50469.1; PID:g1199580
A; Note: the translated sequence in GenBank entry HSU28694, release 113.0,
PIDN:AAC50469.1, differs from the published sequence in having 281-Leu
C:Genetics:
A; Gene: GDB: CMKBR3
A; Cross-references: GDB:579624; OMIM:601268
A; Map position: 3p21-3p21
C; Superfamily: vertebrate rhodopsin
C; Keywords: G protein-coupled receptor; glycoprotein; phosphoprotein; transmembrane
protein
F;36-60/Domain: transmembrane #status predicted <TMl>
F;71-91/Domain: transmembrane #status predicted <TM2>
F;108-129/Domain: transmembrane #status predicted <TM3>
F;147-171/Domain: transmembrane #status predicted <TM4>
F;205-223/Domain: transmembrane #status predicted <TM5>
F;240-261/Domain: transmembrane #status predicted <TM6>
F;288-305/Domain: transmembrane #status predicted <TM7>
F;24-273,106-183/Disulfide bonds: #status predicted
F;345/Binding site: phosphate (Ser) (covalent) (by casein kinase II) #status predicted
 Query Match
                        77.5%; Score 275; DB 2; Length 355; 100.0%; Pred. No. 1.7e-273;
 Best Local Similarity
 Matches 275; Conservative
                               0; Mismatches
                                                 0; Indels
                                                               0; Gaps
       1 MTTSLDTVETFGTTSYYDDVGLLCEKADTRALMAQFVPPLYSLVFTVGLLGNVVVVMILI 60
Ov
         1 MTTSLDTVETFGTTSYYDDVGLLCEKADTRALMAOFVPPLYSLVFTVGLLGNVVVVMILI 60
Db
      61 KYRRLRIMTNIYLLNLAISDLLFLVTLPFWIHYVRGHNWVFGHGMCKLLSGFYHTGLYSE 120
Qу
         Db
      61 KYRRLRIMTNIYLLNLAISDLLFLVTLPFWIHYVRGHNWVFGHGMCKLLSGFYHTGLYSE 120
```

```
121 IFFIILLTIDRYLAIVHAVFALRARTVTFGVITSIVTWGLAVLAALPEFIFYETEELFEE 180
Qу
         Db
      121 IFFIILLTIDRYLAIVHAVFALRARTVTFGVITSIVTWGLAVLAALPEFIFYETEELFEE 180
     181 TLCSALYPEDTVYSWRHFHTLRMTIFCLVLPLLVMAICYTGIIKTLLRCPSKKKYKAIRL 240
Qу
         181 TLCSALYPEDTVYSWRHFHTLRMTIFCLVLPLLVMAICYTGIIKTLLRCPSKKKYKAIRL 240
Db
     241 IFVIMAVFFIFWTPYNVAILLSSYQSILFGNDCER 275
Qу
         Db
     241 IFVIMAVFFIFWTPYNVAILLSSYQSILFGNDCER 275
RESULT
A45177
chemokine (C-C) receptor 1 - human
N; Alternate names: C-C CKR-1; macrophage inflammatory protein-1-alpha receptor
C: Species: Homo sapiens (man)
C;Date: 30-Sep-1993 #sequence_revision 30-Sep-1993 #text_change 13-Aug-1999
C; Accession: A45177; I55671
R; Neote, K.; DiGregorio, D.; Mak, J.Y.; Horuk, R.; Schall, T.J.
Cell 72, 415-425, 1993
A; Title: Molecular cloning, functional expression, and signaling characteristics of a C-C
chemokine receptor.
A; Reference number: A45177; MUID: 93161416
A; Accession: A45177
A:Status: nucleic acid sequence not shown
A; Molecule type: mRNA
A; Residues: 1-355 <NEO>
A;Cross-references: GB:L10918; NID:g292416; PIDN:AAA36543.1; PID:g292417
A; Experimental source: HL60 cells
A; Note: sequence extracted from NCBI backbone (NCBIP:124876)
R; Gao, J.
J. Exp. Med. 177, 1421-1427, 1993
A; Title: Structure and functional expression of the human macrophage inflammatory 1 alpha
(MIP-lalpha) / RANTES receptor.
A; Reference number: I55671; MUID: 93240122
A; Accession: I55671
A; Status: preliminary; translated from GB/EMBL/DDBJ
A; Molecule type: mRNA
A:Residues: 1-355 <RES>
A; Cross-references: GB:L10918; NID:q292416; PIDN:AAA36543.1; PID:q292417
C:Genetics:
A;Gene: GDB:CMKBR1; CMKR-1
A; Cross-references: GDB:138446; OMIM:601159
A; Map position: 3p21-3p21
C; Superfamily: vertebrate rhodopsin
C; Keywords: disulfide bond; G protein-coupled receptor; glycoprotein; phosphoprotein;
transmembrane protein
F;36-60/Domain: transmembrane #status predicted <TM1>
F;71-91/Domain: transmembrane #status predicted <TM2>
F;108-129/Domain: transmembrane #status predicted <TM3>
F;147-171/Domain: transmembrane #status predicted <TM4>
F;205-223/Domain: transmembrane #status predicted <TM5>
F;240-264/Domain: transmembrane #status predicted <TM6>
F;288-305/Domain: transmembrane #status predicted <TM7>
F;5/Binding site: carbohydrate (Asn) (covalent) #status predicted
F;24-273,106-183/Disulfide bonds: #status predicted
F;345/Binding site: phosphate (Ser) (covalent) (by casein kinase II) #status predicted
 Query Match
                        11.5%; Score 41; DB 2; Length 355;
 Best Local Similarity 100.0%; Pred. No. 1.2e-33;
                              0; Mismatches
                                                             0; Gaps
 Matches
         41: Conservative
                                               0: Indels
                                                                        0:
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
Qу
         Db
     115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
```

```
RESULT
149339
macrophage inflammatory protein-1 alpha receptor - mouse
C; Species: Mus musculus (house mouse)
C;Date: 02-Jul-1996 #sequence_revision 02-Jul-1996 #text_change 13-Aug-1999
C; Accession: I49339
R; Gao, J.L.; Murphy, P.M.
J. Biol. Chem. 270, 17494-17501, 1995
A; Title: Cloning and differential tissue-specific expression of three mouse beta
chemokine receptor-like genes, including the gene for a functional macrophage
inflammatory protein-1 alpha receptor.
A; Reference number: I49339; MUID: 95340546
A; Accession: I49339
A; Status: preliminary; translated from GB/EMBL/DDBJ
A; Molecule type: DNA
A; Residues: 1-355 < RES>
A;Cross-references: EMBL:U28404; NID:g881547; PIDN:AAA89153.1; PID:g881548
C; Superfamily: vertebrate rhodopsin
                           9.3%; Score 33; DB 2; Length 355;
  Query Match
  Best Local Similarity 100.0%; Pred. No. 1.9e-25;
                                 0; Mismatches
          33; Conservative
                                                 0; Indels
                                                                 0; Gaps
                                                                             0:
  Matches
      116 GLYSEIFFIILLTIDRYLAIVHAVFALRARTVT 148
Qу
          Db
      116 GLYSEIFFIILLTIDRYLAIVHAVFALRARTVT 148
RESULT
JC2443
chemokine (C-C) receptor 2, splice form B - human
N; Alternate names: C-C CKR-2; monocyte chemoattractant protein 1 receptor; monocyte
chemotactin 1 receptor
C; Species: Homo sapiens (man)
C;Date: 21-Feb-1995 #sequence revision 05-Apr-1995 #text change 20-Jun-2000
C; Accession: JC2443; I38463
R; Yamagami, S.; Tokuda, Y.; Ishii, K.; Tanaka, H.; Endo, N.
Biochem. Biophys. Res. Commun. 202, 1156-1162, 1994
A; Title: cDNA cloning and functional expression of a human monocyte chemoattractant
protein 1 receptor.
A; Reference number: JC2443; MUID: 94324942
A; Accession: JC2443
A; Molecule type: mRNA
A; Residues: 1-360 < YAM>
A;Cross-references: DDBJ:D29984; NID:g531246; PIDN:BAA06253.1; PID:g531247
R; Charo, I.F.; Myers, S.J.; Herman, A.; Franci, C.; Connolly, A.J.; Coughlin, S.R.
Proc. Natl. Acad. Sci. U.S.A. 91, 2752-2756, 1994
A; Title: Molecular cloning and functional expression of two monocyte chemoattractant
protein 1 receptors reveals alternate splicing of the carboxyl-terminal tails.
A; Reference number: A53477; MUID: 94195821
A; Accession: I38463
A; Status: preliminary
A; Molecule type: mRNA
A: Residues: 1-360 < RES>
A; Cross-references: EMBL: U03905; NID: g472557; PIDN: AAA19120.1; PID: g472558
C:Genetics:
A; Gene: GDB: CMKBR2
A; Cross-references: GDB:337364; OMIM:601267
A; Map position: 3p21-3p21
C; Superfamily: vertebrate rhodopsin
C; Keywords: alternative splicing; G protein-coupled receptor; glycoprotein; transmembrane
protein
F;43-70/Domain: transmembrane #status predicted <TMl>
F;81-100/Domain: transmembrane #status predicted <TM2>
F;115-136/Domain: transmembrane #status predicted <TM3>
F;154-178/Domain: transmembrane #status predicted <TM4>
```

```
F;207-226/Domain: transmembrane #status predicted <TM5>
F;244-268/Domain: transmembrane #status predicted <TM6>
F;287-309/Domain: transmembrane #status predicted <TM7>
F;14/Binding site: carbohydrate (Asn) (covalent) #status predicted
F;113-190/Disulfide bonds: #status predicted
                          6.2%; Score 22; DB 2; Length 360;
  Best Local Similarity 100.0%; Pred. No. 3.7e-14;
  Matches 22; Conservative
                               0; Mismatches 0; Indels
                                                                0; Gaps
                                                                            0;
      121 IFFIILLTIDRYLAIVHAVFAL 142
          131111111111
Dh
      128 IFFIILLTIDRYLAIVHAVFAL 149
RESULT
I38450
chemokine (C-C) receptor 2, splice form A - human
N; Alternate names: C-C CKR-2; monocyte chemoattractant protein 1 receptor; monocyte
chemotactin 1 receptor
C; Species: Homo sapiens (man)
C;Date: 16-Feb-1996 #sequence revision 16-Feb-1996 #text change 13-Aug-1999
C; Accession: I38450
R; Charo, I.F.; Myers, S.J.; Herman, A.; Franci, C.; Connolly, A.J.; Coughlin, S.R.
Proc. Natl. Acad. Sci. U.S.A. 91, 2752-2756, 1994
A; Title: Molecular cloning and functional expression of two monocyte chemoattractant
protein 1 receptors reveals alternate splicing of the carboxyl-terminal tails.
A; Reference number: A53477; MUID: 94195821
A; Accession: I38450
A; Status: preliminary
A; Molecule type: mRNA
A; Residues: 1-374 < RES>
A;Cross-references: EMBL:U03882; NID:g472555; PIDN:AAA19119.1; PID:g472556
C:Genetics:
A; Gene: GDB: CMKBR2
A; Cross-references: GDB:337364; OMIM:601267
A; Map position: 3p21-3p21
C; Superfamily: vertebrate rhodopsin
C; Keywords: alternative splicing; G protein-coupled receptor; glycoprotein; transmembrane
F;44-68/Domain: transmembrane #status predicted <TMl>
F;79-99/Domain: transmembrane #status predicted <TM2>
F;115-136/Domain: transmembrane #status predicted <TM3>
F;154-178/Domain: transmembrane #status predicted <TM4>
F;208-226/Domain: transmembrane #status predicted <TM5>
F;244-265/Domain: transmembrane #status predicted <TM6>
F;292-309/Domain: transmembrane #status predicted <TM7>
F;14/Binding site: carbohydrate (Asn) (covalent) #status predicted
F;32-277,113-190/Disulfide bonds: #status predicted
  Query Match 6.2%; Score 22; DB 2; Length 374; Best Local Similarity 100.0%; Pred. No. 3.8e-14;
                              0; Mismatches 0; Indels
  Matches 22; Conservative
                                                                0; Gaps
                                                                            0:
Qу
     121 IFFIILLTIDRYLAIVHAVFAL 142
          1111111111111111111111111
      128 IFFIILLTIDRYLAIVHAVFAL 149
                                   SUMMARIES
Result
               Ouery
        Score Match Length DB ID
                                                         Description
  No.
  355 100.0 355 1 CKR3_HUMAN P51677 homo sapien 64 18.0 355 1 CKR3_MACMU P56483 macaca mula
    1
```

355 1 CKR3_CERAE

P56492 cercopithec

55 15.5

```
11.5
                     355 1 CKR1_HUMAN
                                                         P32246 homo sapien
        41
                              CKR1 MACMU
                                                         P56482 macaca mula
        41
             11.5
                     355 1
                     355 1 CKR1_MOUSE
                                                         P51675 mus musculu
6
        3.3
              9.3
              9.3
                     359 1
                              CKR3_MOUSE
                                                         P51678 mus musculu
        33
                     358 1 CKR3 CAVPO
                                                         09z2i3 cavia porce
8
        29
              8.2
9
        23
              6.5
                     359 1 CKR3 RAT
                                                         054814 rattus norv
                     352 1 CKR5_CERAE
352 1 CKR5_CERTO
10
        22
              6.2
                                                         P56493 cercopithec
11
        22
              6.2
                                                         062743 cercocebus
                                                         P56439 gorilla gor
12
        22
              6.2
                     352 1 CKR5 GORGO
                     352 1 CKR5_HYLLE
                                                         097883 hylobates 1
13
        22
              6.2
14
        22
              6.2
                     352
                          1
                              CKR5 MACMU
                                                         P79436 macaca mula
                     352 1
                              CKR5 PANTR
15
                                                         P56440 pan troglod
        22
              6.2
              6.2
                     352 1
                              CKR5_PAPHA
                                                         P56441 papio hamad
16
        22
                     352 1 CKR5_PONPY
352 1 CKR5_PYGBI
                                                         097881 pongo pygma
17
        22
              6.2
18
        22
              6.2
                                                         097880 pygathrix b
```

```
RESULT 4
CKR1_HUMAN
     CKR1_HUMAN
                    STANDARD:
                                   PRT:
                                          355 AA.
ID
AC
     P32246;
     01-OCT-1993 (Rel. 27, Created)
DT
DT
     01-OCT-1993 (Rel. 27, Last sequence update)
DT
     16-OCT-2001 (Rel. 40, Last annotation update)
     C-C chemokine receptor type 1 (C-C CKR-1) (CC-CKR-1) (CCR-1) (CCR1)
DE
DE
     (Macrophage inflammatory protein-1 alpha receptor) (MIP-lalpha-R)
     (RANTES-R) (HM145) (LD78 receptor).
DE
GN
     CCR1 OR CMKBR1 OR CMKR1.
     Homo sapiens (Human).
os
OC
     Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC
     Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
OX
     NCBI TaxID=9606;
RN
     [1]
     SEQUENCE FROM N.A.
RP
     MEDLINE=93161416; PubMed=7679328;
RX
     Neote K., Digregorio D., Mak J.Y., Horuk R., Schall T.J.;
RΑ
     "Molecular cloning, functional expression, and signaling
RT
RT
     characteristics of a C-C chemokine receptor.";
RL
     Cell 72:415-425(1993).
RN
     [2]
     SEQUENCE FROM N.A.
RP
     MEDLINE=93240122; PubMed=7683036;
RX
     Gao J.-L., Kuhns D., Tiffany H.L., McDermott D., Li X., Francke U.,
RA
RT
     "Structure and functional expression of the human macrophage
RT
     inflammatory protein 1 alpha/RANTES receptor.";
     J. Exp. Med. 177:1421-1427(1993).
RL
RN
     [3]
RP
     SEQUENCE FROM N.A.
RC
     TISSUE=Monocytes;
RX
     MEDLINE=94092629; PubMed=7505609;
RA
     Nomura H., Nielsen B.W., Matsushima K.;
     "Molecular cloning of cDNAs encoding a LD78 receptor and putative
RT
RT
     leukocyte chemotactic peptide receptors.";
RL
     Int. Immunol. 5:1239-1249(1993).
CC
     -!- FUNCTION: RECEPTOR FOR A C-C TYPE CHEMOKINE. BINDS TO MIP-1-ALPHA,
CC
         MIP-1 DELTA, RANTES, AND MCP-3 AND, LESS EFFICIENTLY, TO MIP-1-
         BETA OR MCP-1 AND SUBSEQUENTLY TRANSDUCES A SIGNAL BY INCREASING
CC
CC
         THE INTRACELLULAR CALCIUM IONS LEVEL. RESPONSIBLE FOR AFFECTING
         STEM CELL PROLIFERATION.
CC
     -!- SUBCELLULAR LOCATION: Integral membrane protein.
CC
CC
     -!- TISSUE SPECIFICITY: WIDELY EXPRESSED IN DIFFERENT HEMATOPOIETIC
CC
         CELLS.
CC
     -!- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.
CC
CC
    This SWISS-PROT entry is copyright. It is produced through a collaboration
     between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC
     the European Bioinformatics Institute. There are no restrictions on its
CC
CC
    use by non-profit institutions as long as its content is in no way
     modified and this statement is not removed. Usage by and for commercial
```

modified and this statement is not removed. Usage by and for commercial

```
entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC
     or send an email to license@isb-sib.ch).
CC
DR
     EMBL; L09230; AAA58408.1; -.
     EMBL; L10918; AAA36543.1; -.
DR
     EMBL; D10925; BAA01723.1; -.
DR
     PIR; A45177; A45177.
DR
DR
     GCRDb; GCR_0498; -.
     GCRDb; GCR 0557; -.
DR
DR
     GCRDb; GCR_0573; -.
DR
     MIM; 601159; -.
DR
     InterPro; IPR000276; GPCR Rhodpsn.
DR
     Pfam; PF00001; 7tm 1; 1.
     PRINTS; PR00237; GPCRRHODOPSN.
DR
     PROSITE; PS00237; G PROTEIN RECEP F1 1; 1.
DR
     PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
DR
KW
     G-protein coupled receptor; Transmembrane; Glycoprotein.
FT
     DOMAIN
                                  EXTRACELLULAR (POTENTIAL).
                         34
                   1
FT
     TRANSMEM
                  35
                         60
                                  1 (POTENTIAL).
     DOMAIN
FT
                  61
                         64
                                  CYTOPLASMIC (POTENTIAL).
     TRANSMEM
FT
                  65
                         91
                                  2 (POTENTIAL).
FT
     DOMAIN
                  92
                        107
                                  EXTRACELLULAR (POTENTIAL).
                                  3 (POTENTIAL).
     TRANSMEM
                 108
FT
                        129
FT
     DOMAIN
                 130
                        146
                                  CYTOPLASMIC (POTENTIAL).
FT
     TRANSMEM
                                  4 (POTENTIAL).
                 147
                        171
FT
     DOMAIN
                 172
                        197
                                  EXTRACELLULAR (POTENTIAL).
     TRANSMEM
FT
                 198
                                  5 (POTENTIAL).
                        223
FT
     DOMAIN
                 224
                        239
                                  CYTOPLASMIC (POTENTIAL).
FT
     TRANSMEM
                 240
                                  6 (POTENTIAL).
                        264
FT
     DOMAIN
                 265
                        281
                                  EXTRACELLULAR (POTENTIAL).
FT
     TRANSMEM
                 282
                        305
                                  7 (POTENTIAL).
                                  CYTOPLASMIC (POTENTIAL).
FT
     DOMAIN
                 306
                        355
     CARBOHYD
                                  N-LINKED (GLCNAC. . .) (POTENTIAL) .
FT
                  5
                          5
FT
                 106
     DISULFID
                        183
                                  BY SIMILARITY.
FT
     CONFLICT
                 337
                        337
                                  E \rightarrow D (IN REF. 3).
SQ
     SEQUENCE
                355 AA; 41172 MW; B2C100FFED275985 CRC64;
  Query Match
                          11.5%; Score 41; DB 1; Length 355;
  Best Local Similarity
                          100.0%; Pred. No. 3.2e-31;
                                 0; Mismatches
  Matches
           41; Conservative
                                                    0; Indels
                                                                  0; Gaps
Qу
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
          Db
      115 TGLYSEIFFIILLTIDRYLAIVHAVFALRARTVTFGVITSI 155
RESULT
CKR1_MOUSE
ID
     CKR1 MOUSE
                    STANDARD:
                                    PRT:
                                           355 AA.
AC
     P51675;
     01-OCT-1996 (Rel. 34, Created)
01-OCT-1996 (Rel. 34, Last sequence update)
DТ
DT
DT
     30-MAY-2000 (Rel. 39, Last annotation update)
DE
     C-C chemokine receptor type 1 (C-C CKR-1) (CC-CKR-1) (CCR-1) (CCR1)
ĎΕ
     (Macrophage inflammatory protein-1 alpha receptor) (MIP-1alpha-R)
DE
     (RANTES-R).
GN
     CCR1 OR CMKBR1.
os
     Mus musculus (Mouse).
OC
     Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC.
     Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.
OX
     NCBI_TaxID=10090;
RN
     [1]
     SEQUENCE FROM N.A.
RΡ
RC
     STRAIN=129/SV; TISSUE=Peritoneal macrophage;
RX
     MEDLINE=96072806; PubMed=7594543;
RA
     Post T.W., Bozic C.R., Rothenberg M.E., Luster A.D., Gerard N.,
RA
     Gerard C.;
RТ
     "Molecular characterization of two murine eosinophil beta chemokine
```

```
RT
    receptors.";
    J. Immunol. 155:5299-5305(1995).
RL
RN
     [2]
     SEQUENCE FROM N.A.
RΡ
    STRAIN=129/SVJ;
RC
    MEDLINE=95340546; PubMed=7542241;
RX
    Gao J.-L., Murphy P.M.;
RA
RT
     "Cloning and differential tissue-specific expression of three mouse
    beta chemokine receptor-like genes, including the gene for a
RT
RТ
     functional macrophage inflammatory protein-1 alpha receptor.";
RL
     J. Biol. Chem. 270:17494-17501(1995).
     -!- FUNCTION: RECEPTOR FOR A C-C TYPE CHEMOKINE. BINDS TO MIP-1-ALPHA,
CC
CC
         RANTES, AND LESS EFFICIENTLY, TO MIP-1-BETA OR MCP-1 AND
CC
         SUBSEQUENTLY TRANSDUCES A SIGNAL BY INCREASING THE INTRACELLULAR
CC
         CALCIUM IONS LEVEL. RESPONSIBLE FOR AFFECTING STEM CELL
CC
         PROLIFERATION.
CC
     -!- SUBCELLULAR LOCATION: Integral membrane protein.
CC
     -!- TISSUE SPECIFICITY: DETECTED IN THE HEART, SPLEEN, LUNG,
        PERITONEAL EXUDATE CELLS AND LEUKOCYTES.
CC
     -!- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.
CC
CC
     ______
CC
    This SWISS-PROT entry is copyright. It is produced through a collaboration
CC
    between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC
    the European Bioinformatics Institute. There are no restrictions on its
CC
    use by non-profit institutions as long as its content is in no way
    modified and this statement is not removed. Usage by and for commercial
CC
CC
    entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC
    or send an email to license@isb-sib.ch).
CC
    EMBL; U29678; AAA86119.1; -.
DR
    EMBL; U28404; AAA89153.1; -.
DR
DR
    GCRDb; GCR 1672; -.
DR
    GCRDb; GCR 1698; -.
DR
    MGD; MGI:104618; Cmkbr1.
    InterPro; IPR000276; GPCR Rhodpsn.
DR
    Pfam; PF00001; 7tm 1; 1.
DR
DR
    PRINTS; PR00237; GPCRRHODOPSN.
    PROSITE; PS00237; G PROTEIN RECEP F1 1; 1.
DR
DR
    PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
    G-protein coupled receptor; Transmembrane.
KW
FT
    DOMAIN
                                 EXTRACELLULAR (POTENTIAL).
                 1
                       34
    TRANSMEM
FT
                 35
                        60
                                 1 (POTENTIAL).
FT
    DOMAIN
                 61
                        64
                                 CYTOPLASMIC (POTENTIAL).
    TRANSMEM
FT
                 65
                        91.
                                 2 (POTENTIAL).
FT
    DOMAIN
                 92
                       107
                                 EXTRACELLULAR (POTENTIAL).
    TRANSMEM
FT
                108
                       129
                                 3 (POTENTIAL).
FT
    DOMAIN
                130
                                 CYTOPLASMIC (POTENTIAL).
                       146
    TRANSMEM
FT
                147
                       171
                                 4 (POTENTIAL).
FT
    DOMAIN
                172
                       197
                                EXTRACELLULAR (POTENTIAL).
FT
    TRANSMEM
                198
                                 5 (POTENTIAL).
                       223
FT
    DOMAIN
                224
                       239
                                 CYTOPLASMIC (POTENTIAL).
FT
    TRANSMEM
                240
                                 6 (POTENTIAL).
                       264
    DOMAIN
FT
                265
                       281
                                 EXTRACELLULAR (POTENTIAL).
                282
FT
    TRANSMEM
                       305
                                 7 (POTENTIAL).
FT
    DOMAIN
                306
                       355
                                 CYTOPLASMIC (POTENTIAL).
FT
    DISULFID
                106
                       183
                                 BY SIMILARITY.
FT
    CONFLICT
                55
                        55
                                 M \rightarrow V (IN REF. 2).
               355 AA; 40901 MW; FCE9FFF70E6F38B1 CRC64;
so
    SEQUENCE
 Query Match 9.3%; Score 33; DB 1; Length 355; Best Local Similarity 100.0%; Pred. No. 1.3e-23;
          33; Conservative
                                0; Mismatches
                                                                0; Gaps
                                                                            0;
Qу
     116 GLYSEIFFIILLTIDRYLAIVHAVFALRARTVT 148
          1111111111111111
```

Db

116 GLYSEIFFIILLTIDRYLAIVHAVFALRARTVT 148